

STELLAR POPULATIONS OF THE DWARF GALAXY UKS 2323–326 IN THE SCULPTOR GROUP

MYUNG GYOON LEE

Department of Astronomy, Seoul National University, Seoul 151-742, Korea
 Electronic mail: mglee@astrog.snu.ac.kr

AND

YONG-IK BYUN

Department of Astronomy and Center for Space Astrophysics, Yonsei University, Seoul 120-749, Korea
 Electronic mail: byun@darksky.yonsei.ac.kr

Draft version February 1, 2008

ABSTRACT

We present deep *BVRI* CCD photometry of the stars in the dwarf irregular galaxy UKS 2323–326 in the Sculptor Group. The color-magnitude diagrams of the measured stars in UKS 2323–326 show a blue plume which consists mostly of young stellar populations, and a well-defined red giant branch (RGB). The tip of the RGB is found to be at $I_{\text{TRGB}} = 22.65 \pm 0.10$ mag. From this the distance to this galaxy is estimated to be $d = 2.08 \pm 0.12$ Mpc. The corresponding distance of this galaxy from the center of the Local Group is 1.92 Mpc, showing that it is located outside the Local Group. The large distance, combined with its velocity information, indicates that it is very likely to be a member of the Sculptor Group. The mean metallicity of the red giant branch is estimated to be $[\text{Fe}/\text{H}] = -1.98 \pm 0.17$ dex. Total magnitudes of UKS 2323–326 ($< r_H = 70$ arcsec) are derived to be $B^T = 14.07$ mag, $V^T = 13.50$ mag, $R^T = 13.18$ mag, and $I^T = 12.83$ mag, and the corresponding absolute magnitudes are $M_B = -12.58$ mag, $M_V = -13.14$ mag, $M_R = -13.45$ mag, and $M_I = -13.79$ mag. Surface brightness profiles of the central part of UKS 2323–326 are approximately consistent with a King model with a core concentration parameter $c = \log(r_t/r_c) \approx 1.0$, and those of the outer part follow an exponential law with a scale length of 21 arcsec. The magnitudes and colors of the brightest blue and red stars in UKS 2323–326 (BSG and RSG) are measured to be, respectively, $< V(3) >_{BSG} = 20.33 \pm 0.25$ mag, $< (B-V)(3) >_{BSG} = 0.14 \pm 0.07$ mag, $< V(3) >_{RSG} = 20.74 \pm 0.18$ mag, and $< (B-V)(3) >_{RSG} = 1.35 \pm 0.08$ mag. The corresponding absolute magnitudes are derived to be $< M_V(3) >_{BSG} = -6.31$ mag and $< M_V(3) >_{RSG} = -5.91$ mag, which are about half magnitude fainter than those expected from conventional correlations with galaxy luminosity.

[To appear in the *Astronomical Journal* in August, 1999]

Subject headings: galaxy: evolution — galaxies: irregular — galaxies: individual (UKS 2323–326) — galaxies: stellar content — galaxies: photometry — Distance scale

1. INTRODUCTION

UKS 2323–326 (UGCA 438) is a faint dwarf irregular galaxy in Sculptor discovered by Longmore et al. (1978). In the discovery paper, Longmore et al. estimated the distance to this galaxy from the rough estimate of the magnitude of the brightest stars, to be 1.3 Mpc with a large error of ± 50 %. Since then this galaxy has been often considered to be a member of the Local Group (Longmore et al. 1978, Mateo 1998). On the other hand, the velocity of UKS 2323–326 from the center of the Local Group, 82 km s^{-1} (Longmore et al. 1978), suggests that it may not belong to the Local Group (van den Bergh 1994, Lee 1995a). Accurate measurement of the distance to this galaxy is needed to resolve this problem.

Longmore et al. (1978) measured the HI flux of $15 \pm 3 \text{ Jy km s}^{-1}$ at the heliocentric velocity of $62 \pm 5 \text{ km s}^{-1}$, and derived, using eye-estimate with the photographic plates, the magnitude of the brightest blue stars in UKS 2323–326, $B = 19.3 \pm 0.5$ mag. After the discovery paper, however, there has been published no detailed photometric study for this galaxy.

In this paper we present a study of stellar populations of UKS 2323–326 based on deep *BVRI* CCD photometry,

and show that this galaxy is outside the Local Group and is a member of the Sculptor Group. This paper is composed as follows. Section 2 describes the observations and data reduction, and Section 3 investigates the morphological structure of UKS 2323–326. Section 4 presents the color-magnitude diagrams of UKS 2323–326, Section 5 estimates the distance to UKS 2323–326. Section 6 presents the surface photometry of UKS 2323–326 and Section 7 discusses the group membership, the stellar populations, and the brightest stars of UKS 2323–326. Finally, summary and conclusion are given in Section 8.

2. OBSERVATIONS AND DATA REDUCTION

BVRI CCD images of UKS 2323–326 were obtained on the photometric night of 1994 October 7 (UT) using the University of Hawaii 2.2 m telescope at Mauna Kea. Table 1 lists the journal of the observations of UKS 2323–326. A grey scale map of the *V*-band CCD image of UKS 2323–326 is displayed in Fig. 1. The size of the field of view is $7'.5 \times 7'.5$ and the (2×2 binned) pixel scale of the CCD is $0.44 \text{ arcsec pixel}^{-1}$.

For the analysis of the data we have divided the field covered by our CCD images into three regions as shown in Fig. 1: the C-region which covers the central region

($r < 44''$) of UKS 2323–326, the I-region which covers the outer region ($44'' < r < 77''$) of the galaxy, and the F-region which represents a control field with the same area of the C-region plus I-region.

Instrumental magnitudes of the stars in the CCD images were derived using DoPHOT (Schechter, Saha, & Mateo 1993). These magnitudes were transformed onto the standard system using the standard stars observed during two nights including the same night (Landolt 1992). The transformation equations we derived from the photometry of the standard stars are: $V = v - 0.075(b - v) - 0.118X + \text{constant}$, $(B - V) = 1.130(b - v) - 0.111X + \text{constant}$, $(V - R) = 0.970(v - r) - 0.030X + \text{constant}$, and $I = i + 0.050(v - i) - 0.082X + \text{constant}$, where upper cases and lower cases represent, respectively, the standard system and instrumental system. X represents the air mass. The rms scatter of the solutions are $0.01 - 0.02$ mag. The total number of stars which were measured at V and at least one other color in the CCD image is $\sim 2,000$. Table 2 lists $BVRI$ photometry of the measured bright stars with $V < 22.5$ mag in the field of UKS 2323–326. The coordinates X and Y in Table 2 are given in units of pixel ($= 0.44$ arcsec), and increase to the east and to the south, respectively.

3. MORPHOLOGICAL STRUCTURE

Unfortunately there is a very bright star $20''$ south-east of the center of UKS 2323–326, as shown in Fig. 1. It is about 14 mag foreground star (Longmore et al. 1978) and is saturated in all our CCD images. Fig. 1 shows that there are several tens of bright stars concentrated in the central region of UKS 2323–326. These stars are considered mostly to be the members of UKS 2323–326 and are young massive stars as shown later. The main body of the galaxy is mostly seen inside the C-region, and is slightly elongated along the south-east direction. However, the outer part of the galaxy seen at the faint level extends out to the boundary of the I-region ($r = 77''$), and is impressively almost circular. There are few bright stars in the I-region. Instead there are many faint stars seen better in R and I images of the I-region, which are probably old red giant stars as seen later. Therefore UKS 2323–326 is a circular galaxy, in the central region of which there are irregularly distributed some bright stars. This kind of structure that some young stars are irregularly distributed against the smooth background of old stellar populations is common among irregular galaxies (Sandage 1971, Lee 1993, Lee, Freedman, & Madore 1993, Minniti & Zijlstra 1996).

4. COLOR-MAGNITUDE DIAGRAMS

UKS 2323–326 is located 71 deg below the galactic plane in the sky so that the foreground reddening for UKS 2323–326 is expected to be very low. We adopt in this study the foreground reddening value of $E(B - V) = 0.014$ mag for UKS 2323–326 given by Schlegel, Finkbeiner, & Davis (1998).

We display $V-(B - V)$ and $I-(V - I)$ diagrams of 520 measured stars in the C-region, 280 measured stars in the I-region, and 89 measured stars in the F-region in Figs. 2 and 3. Note that the area of the (C+I)-region in the field is the same as that of the F-region so that we can estimate

the contamination due to foreground stars by comparing directly the diagrams of each region.

Several distinguishable features of the stars in UKS 2323–326 are seen in Figs. 2 and 3. First, there is a blue plume of bright stars with $(B - V) < 0.4$ in the C-region (filled circles). Comparison of the C-region and the F-region in Fig. 2 shows that these bright blue stars are mostly members of UKS 2323–326. The brightest end of the blue plume extends up to $V \approx 20.0$ mag and $(B - V) \approx -0.1$. These stars are mostly massive stars which were formed recently. Stars in the I-region are all fainter than $V = 23$ mag (open circles), much fainter than those bright stars in the C-region. This shows that there was little star formation in the outer part of UKS 2323–326 recently.

Secondly, there are five red bright stars with $20.0 < V < 21.5$ mag and $1.2 < (B - V) < 1.5$ in the C-region in Fig. 2. This area of the F-region color-magnitude diagram is devoid of stars. Therefore these stars are probably the members of UKS 2323–326. These stars are considered to be red supergiant stars which were formed recently.

Thirdly, Fig. 3 shows that there is a strong concentration of red stars fainter than $I \approx 21.5$ mag in the (C+I)-region. Most of these stars are probably old red giant branch (RGB) stars of UKS 2323–326.

5. DISTANCE AND METALLICITY

We estimate the distance to UKS 2323–326 using the I magnitude of the tip of the RGB (TRGB), as described in Da Costa & Armandroff (1990) and Lee, Freedman, & Madore (1993). The I magnitude of the TRGB is estimated using the $I - (V - I)$ diagram in Fig. 3 and the luminosity function of red giant stars. Fig. 4 shows the I -band luminosity function of the measured red stars in the I-region (solid line) and the C-region (dotted line), from which the contribution due to field stars was subtracted. We also plot the luminosity functions of the similar stars in the field with the same area as the (C+I) region in Fig. 4, which shows that the contribution due to field stars is very small.

The I-region is more appropriate for measuring the magnitude of the TRGB than the C-region, because there are several young bright stars in the C-region and because the crowding is more severe in the C-region than in the I-region. In Fig. 4, as the magnitude increases, there is a sudden increase at $I = 22.65 \pm 0.10$ mag in the luminosity function of the I-region, which corresponds to the TRGB seen in the color-magnitude diagram in Fig. 3. Several stars brighter than the TRGB are mostly asymptotic giant stars of intermediate age, also seen in other dwarf galaxies (Lee 1995b). A dip seen at the magnitude of the TRGB in the luminosity function of the C-region is a statistical fluke.

The mean color of the TRGB is estimated to be $(V - I) = 1.43 \pm 0.05$. The bolometric magnitude of the TRGB is then calculated from $M_{\text{bol}} = -0.19[\text{Fe}/\text{H}] - 3.81$. Adopting a mean metallicity of $[\text{Fe}/\text{H}] = -1.98 \pm 0.17$ dex as estimated below, we obtain a value for the bolometric magnitude of $M_{\text{bol}} = -3.43$ mag. The bolometric correction at I for the TRGB is estimated to be $\text{BC}_I = 0.54$ mag, adopting a formula for the bolometric correction $\text{BC}_I = 0.881 - 0.243(V - I)_{\text{TRGB}}$. The intrinsic I magnitude of the TRGB is then given by $M_I = M_{\text{bol}} - \text{BC}_I = -3.97$

mag. Finally the distance modulus of UKS 2323–326 is obtained: $(m - M)_0 = 26.59 \pm 0.12$ mag (corresponding to a distance of 2.08 ± 0.12 Mpc) for an adopted extinction of $A_I = 0.03$ mag.

We have estimated the mean metallicity of the RGB stars in UKS 2323–326 using the $(V - I)$ color of the stars 0.5 mag fainter than the TRGB, $(V - I)_{-3.5}$. This color is measured from the median value of the colors of 17 red giant branch stars with $I = 23.10 \pm 0.10$ mag, to be $(V - I)_{-3.5} = 1.29 \pm 0.04$. From this value we estimate the mean metallicity to be $[\text{Fe}/\text{H}] = -1.98 \pm 0.17$ dex. In Fig. 5 we overlaid the loci of the red giant branches of Galactic globular clusters, of M15, M2, and NGC 1851, shifted according to the distance and reddening of UKS 2323–326. The metallicities of M15, M2 and NGC 1851 are $[\text{Fe}/\text{H}] = -2.17, -1.58$ and -1.29 dex, respectively. Fig. 5 shows that the bright part of the RGB of UKS 2323–326 is located well between those of M15 and M2. The broadening of the faint part of the RGB is mostly due to the photometric errors, as shown by the error bars in Fig. 5. Thus the mean metallicity of the RGB stars in UKS 2323–326 is very low, and is close to the lowest end in the metallicity of dwarf irregular galaxies (Lee, Freedman, & Madore 1993; Lee 1995a,b; Mateo 1998; Lee et al. 1999).

6. SURFACE PHOTOMETRY

It is very difficult to derive reliably the surface photometry of UKS 2323–326, because of the presence of very bright foreground star close to the center of the galaxy. We have obtained the surface photometry of UKS 2323–326 as follows. First, we removed in the original CCD images the images of very bright foreground star and several other bright stars which were obviously considered to be foreground stars in the area of UKS 2323–326, using IMEDIT in IRAF. Then we performed aperture photometry of UKS 2323–326 using the circular annular aperture. The value of the sky background was estimated from the mean intensity of the F-region. Subtraction of the very bright 14 mag star from the image of the galaxy is very difficult. The error due to this problem is not easy to quantify, but may be significant in the photometry of the galaxy. Therefore the surface photometry of the galaxy derived in this study is only approximate.

The results of surface photometry of UKS 2323–326 are listed in Table 3 and are displayed in Fig. 6. In Table 3 r_{eff} presents the mean major radius of an annular aperture, and r_{out} represents the outer radius of an annular aperture. Fig. 6(a) shows that the surface brightness profiles are almost flat in the central region of the galaxy ($r < 20$ arcsec), and follow approximately the exponential law in the outer part ($r > 20$ arcsec). We fit the V surface brightness profile of the inner region of UKS 2323–326 with the single-mass isotropic King models (King 1966) and that of the outer region with the exponential law in Fig. 7. Fig. 7 shows that the surface brightness profile of the inner region is roughly fit by a King model with a core concentration parameter $c = \log(r_t/r_c) \approx 1.0$, where r_c and r_t represent the core radius and tidal radius, respectively. The surface brightness profiles of the outer region are fit roughly by an exponential law with a scale length of $r = 21$ arcsec = 210 pc. Fig. 6(b) illustrates the surface color profiles for the region at $r < 40$ arcsec. The colors files are plotted only for the region at $r < 40$ arcsec, beyond which the errors of

the colors are too large. The color profiles are almost constant and become slowly redder outward from the center to $r \approx 40$ arcsec. The rapid change of the color profiles right close to the center is due to a bright star in the central region. The properties of the surface brightness and color profiles of UKS2323–326 are similar to those of other dwarf galaxies (Kormendy & Djorgovski 1989, Mateo 1998).

From the data of the surface photometry we have derived several basic parameters of UKS 2323–326 as follows. The standard radius and Holmberg radius of UKS 2323–326 are measured to be $r_{25} = 46$ arcsec = 460 pc and $r_H = 70$ arcsec = 700 pc, respectively. These radii are similar, respectively, to the radii of the C-region and I-region shown in Fig. 1. Note that our value for the Holmberg radius is much larger than the value Longmore et al. (1978) derived roughly from their photographic plate, $r_H = 45$ arcsec.

The central surface brightness of UKS 2323–326 is measured to be $\mu_B(0) = 23.4$ mag arcsec $^{-2}$, $\mu_V(0) = 22.9$ mag arcsec $^{-2}$, $\mu_R(0) = 22.6$ mag arcsec $^{-2}$, and $\mu_I(0) = 22.2$ mag arcsec $^{-2}$. The total magnitudes of UKS 2323–326 within r_H are derived to be $B^T = 14.07$ mag, $V^T = 13.50$ mag, $R^T = 13.18$ mag, and $I^T = 12.83$ mag, and the corresponding absolute magnitudes are $M_B = -12.58$ mag, $M_V = -13.14$ mag, $M_R = -13.45$ mag, and $M_I = -13.79$ mag. B total magnitude of UKS 2323–326 derived in this study, $B^T = 14.07$ mag, is about 1 mag brighter than that given by Longmore et al. (1978), $B^T = 15.2$ mag.

7. DISCUSSION

7.1. The Group Membership of UKS 2323–326

We have measured the distance to UKS 2323–326 to be $d = 2.08 \pm 0.12$ Mpc from the I -band magnitude of the TRGB, which is an accurate distance indicator for resolved galaxies (Lee, Freedman, & Madore 1993, Salaris & Cassisi 1998). Our value is 1.6 times larger and much more accurate than the value Longmore et al. (1978) suggested, from the eye-estimate of the magnitude of the blue brightest stars, $d = 1.3$ Mpc with an error of ± 50 %. Our result shows that UKS 2323–326 is obviously outside the Local Group. The mean distance of the Sculptor group is 2.5 Mpc and the radius of the Sculptor group is ~ 1.1 Mpc (Puche & Carignan 1988, Côte et al. 1997, Jerjen, Freeman, & Binggeli 1998). UKS2323–326 is located in the western boundary of the Sculptor group in the sky (see the map of the Sculptor group members given in Fig. 4 of Côte et al. 1997). Therefore UKS 2323–326 is definitely considered to be a member of the Sculptor group.

This conclusion is consistent with the measured velocity of UKS 2323–326. From the measured heliocentric velocity of this galaxy, $v = 62 \pm 5$ km s $^{-1}$, the velocity of UKS 2323–326 from the center of the Local Group is calculated to be 84 ± 5 km s $^{-1}$ (Longmore et al. 1978, Lee 1995b). This value is 24 km s $^{-1}$ larger than the maximum value of 60 km s $^{-1}$ typically used as the boundary of the Local Group, showing that this galaxy is probably outside the boundary of the Local Group.

Côte et al. (1997) excluded UKS 2323–326 in the list of dwarf candidates for the Sculptor group, based on the results given by Longmore et al. (1978). However, our results show that this galaxy should be included as a member of the Sculptor group.

7.2. Stellar Populations in UKS 2323–326

In the discovery paper Longmore et al. (1978) detected no HII region in UKS 2323–326 (which was confirmed later by Miller 1996) and pointed out that the low optical surface brightness and blue color of this galaxy is difficult to reconcile using current evolutionary models of late-type galaxies. However, it is shown here that the blue color of this galaxy is due to a presence of young stellar populations.

We have investigated roughly the properties of young stellar populations seen in Fig. 2 using the theoretical isochrones. In Fig. 8, we overlay, in the $V - (B - V)$ diagram of UKS 2323–326, three isochrones for the metallicity of $Z = 0.001$ ($[\text{Fe}/\text{H}] = -1.2$ dex) and ages of 10, 30, and 100 Myrs given by the Padova Group (Bertelli et al. 1994). Fig. 8 shows (1) that the brightest blue and red stars in UKS 2323–326 match roughly the blue and red loops of the isochrone with an age of 30 Myrs, and (2) that the brightest main-sequence at $V \approx 21.5$ (about 1 mag below the brightest blue supergiants) indicates an age of 10 Myrs. These results show that stars were formed as recently as 10 Myrs ago in UKS 2323–326. On the other hand, a well-developed RGB shown in the $I - (V - I)$ diagram of Fig. 3 shows that the bulk of the stars in UKS 2323–326 were formed before a few Gyrs ago, and a small number of AGB stars above the TRGB were probably formed a few Gyrs ago.

Fig. 9 displays the V luminosity function of the main-sequence stars with $(B - V) < 0.4$ in UKS 2323–326. The luminosity function we derived is incomplete for the faint end, but is reasonably complete for $V < 23$ mag. The bright part of the luminosity function ($21.4 < V < 23$ mag, $-4.2 < M_V < -3.6$ mag) is approximately fit by a line with a logarithmic slope of 0.52 ± 0.12 , which is similar to those of other irregular and spiral galaxies (Freedman 1986, Hoessel 1986).

The integrated HI flux of this galaxy was measured to be $15 \pm 3 \text{ Jy km s}^{-1}$ (Longmore et al. 1978, Longmore et al. 1982), and Longmore et al. (1978) pointed out that the ratio of the HI mass to B luminosity of this galaxy they derived, $M_{\text{HI}}/L_B = 2.7 M_{\odot}/L_{\odot}$, is rather large for normal irregular galaxies. With better data for the distance and luminosity obtained in this study, we derive an HI mass, $M_{\text{HI}} = 1.5 \times 10^7 M_{\odot}$, a B luminosity, $L_B = 1.67 \times 10^7 L_{\odot}$, and a ratio of $M_{\text{HI}}/L_B = 0.9 M_{\odot}/L_{\odot}$ which is much lower than the value given by Longmore et al. (1978). Our value for the ratio of the HI mass to B luminosity is typical for dwarf irregular galaxies (Mateo 1998).

7.3. The Brightest Blue and Red Stars in UKS 2323–326

Longmore et al. (1978) estimated roughly from the survey plate the magnitude of the brightest blue stars in UKS 2323–326 to be $B = 19.3 \pm 0.5$ mag, and used this result to derive the distance to this galaxy, obtaining a value of 1.3 Mpc. With our photometry we can investigate in detail the properties of the brightest stars in this galaxy.

In the $V - (B - V)$ diagram of the C-region of UKS 2323–326 (filled circles) shown in Fig. 2, it is obvious which stars are the three brightest blue and red stars in UKS 2323–326. Three brightest blue stars (called BSG) are IDs 1055, 1002, and 1116, and three brightest red stars (called RSG) are IDs 956, 1058, and 821, as listed in Table

2. There is one very blue star (ID 1140) as bright as ID 1116, but the color is too blue, $(B - V) = -0.49$ to be a normal star. Inspection of the image of this object shows that the image is slightly asymmetric so that the point-spread function fitting photometry of this object may not be as good as that of other normal stars. So we did not include it in the sample of the three brightest blue stars. The mean magnitudes and colors of these three brightest blue and red stars in UKS 2323–326 are derived to be, respectively, $\langle V(3) \rangle_{\text{BSG}} = 20.33 \pm 0.25$ mag and $\langle (B - V)(3) \rangle_{\text{BSG}} = 0.14 \pm 0.07$, and $\langle V(3) \rangle_{\text{RSG}} = 20.74 \pm 0.18$ mag and $\langle (B - V)(3) \rangle_{\text{RSG}} = 1.35 \pm 0.08$. (If ID 1116 is replaced by ID 1140, the mean values will be slightly changed: $\langle V(3) \rangle_{\text{BSG}} = 20.32 \pm 0.28$ mag and $\langle (B - V)(3) \rangle_{\text{BSG}} = -0.04 \pm 0.32$.) The corresponding absolute magnitudes and colors are $\langle M_V(3) \rangle_{\text{BSG}} = -6.31 \pm 0.25$ mag, $\langle (B - V)(3) \rangle_{\text{BSG},0} = 0.14 \pm 0.07$, $\langle M_V(3) \rangle_{\text{RSG}} = -5.91 \pm 0.18$ mag, and $\langle (B - V)(3) \rangle_{\text{RSG},0} = 1.34 \pm 0.08$, respectively.

The luminosity of the brightest stars in galaxies is known to be correlated with the luminosity of the parent galaxies. Recently Lyo & Lee (1997) presented, from the analysis of 17 galaxies (with $M_B < -14$ mag) to which Cepheid distances are available, calibrations for the relation between the magnitudes of the brightest stars and the magnitudes of the parent galaxies: $\langle M_V(3) \rangle_{\text{BSG}} = 0.30 M_B(\text{gal}) - 3.02$ with $\sigma = 0.55$ mag, and $\langle M_V(3) \rangle_{\text{RSG}} = 0.21 M_B(\text{gal}) - 3.84$ with $\sigma = 0.47$ mag. Using these relations, we derive $\langle M_V(3) \rangle_{\text{BSG}} = -6.79$ mag and $\langle M_V(3) \rangle_{\text{RSG}} = -6.48$ mag for the absolute magnitude of UKS 2323–326 as derived in the previous section. Thus the magnitudes of the brightest stars in UKS 2323–326 are 0.5–0.6 mag fainter than those expected from the relation for the bright galaxies.

8. SUMMARY AND CONCLUSION

We have presented a study of the stellar populations in the dwarf irregular galaxy UKS 2323–326 based on deep $BVRI$ CCD photometry. The primary results obtained in this study are summarized as follows and the basic information of UKS 2323–326 is listed in Table 4.

(1) $BVRI$ color-magnitude diagrams of the stars in the $7'.35 \times 7'.35$ area of UKS 2323–326 have been presented. These color-magnitude diagrams exhibit a blue plume, a well-defined RGB, and a small number of AGB stars with intermediate age.

(2) The tip of the RGB is found to be at $I = 22.65 \pm 0.10$ mag and $(V - I) = 1.43 \pm 0.05$ mag. From this value we derive a distance modulus of UKS 2323–326 of $(m - M)_0 = 26.59 \pm 0.12$ mag, and a distance of 2.08 ± 0.12 Mpc. The corresponding distance of UKS 2323–326 from the center of the Local Group is derived to be 1.92 Mpc. From this result and the systemic velocity of UKS 2323–326 ($v = 62 \text{ km s}^{-1}$), we conclude that UKS 2323–326 is outside the Local Group and is a member of the Sculptor group.

(3) The mean color of the RGB at $M_I = -3.5$ mag is $(V - I) = 1.29 \pm 0.04$ mag. From this value we obtain a mean metallicity of the RGB: $[\text{Fe}/\text{H}] = -1.98 \pm 0.17$ dex. The metallicity of the RGB in UKS2323–326 is close to the lowest end in the metallicity of dwarf irregular galaxies.

(4) The total magnitudes of UKS 2323–326 within r_H are derived to be $M_B = -12.58$ mag, $M_V = -13.14$ mag, $M_R = -13.45$ mag, and $M_I = -13.79$ mag. The cen-

tral surface brightness is measured to be $\mu_B(0) = 23.4$ mag arcsec $^{-2}$ and $\mu_V(0) = 22.9$ mag arcsec $^{-2}$. Surface brightness profiles of the central part of UKS 2323–326 are approximately consistent with a King model with a core concentration parameter $c = \log(r_t/r_c) \approx 1.0$, and those of the outer part follow an exponential law.

(5) The magnitudes of three brightest blue and red stars in UKS 2323–326 are derived: $\langle M_V(3) \rangle_{BSG} = -6.31 \pm 0.25$ mag and $\langle M_V(3) \rangle_{RSG} = -5.91 \pm 0.18$ mag, which are about half magnitude fainter than those expected from conventional correlations with galaxy lumi-

nosity.

The authors are grateful to Eunhyeuk Kim for providing the surface photometry program, and to Sang Chul Kim for reading the manuscript carefully. This research is supported by the Ministry of Education, Basic Science Research Institute grant No.BSRI-98-5411 (to M.G.L.) and by Creative Research Initiatives Program of the Korean Ministry of Science and Technology and also by Yonsei University Research Grant (to Y.I.B.).

REFERENCES

- Bertelli, G., Bressan, A., Fagotto, F., Chiosi, C., & Nasi, E. 1994, *A&AS*, 106, 275
 Côte, S., Freeman, K. C., Carignan, C., & Quinn, P. J. 1997, *AJ*, 114, 1313
 Da Costa, G. S., & Armandroff, T. E. 1990, *AJ*, 100, 162
 Freedman, W. L. 1986, *Luminous Stars and Associations in Galaxies*, edited by C. W. H. de Loore, A. J. Willis, & P. Laskarides, IAU Symposium No. 116 (Reidel, Dordrecht), p. 61
 Hoessel, J. 1986, *Luminous Stars and Associations in Galaxies*, edited by C. W. H. de Loore, A. J. Willis, & P. Laskarides, IAU Symposium No. 116 (Reidel, Dordrecht), p. 439
 Jerjen, H., Freeman, K. C., & Binggeli, B. 1998, *AJ*, 116, 2873
 King, I. 1966, *AJ*, 71, 276
 Kormendy, J., & Djorgovski, S. 1989, *ARA&A*, 27, 235
 Landolt, A. U. 1992, *AJ*, 104, 340
 Lee, M. G. 1993, *ApJ*, 408, 409
 Lee, M. G., Freedman, W. L., & Madore, B. F. 1993, *ApJ*, 417, 553
 Lee, M. G. 1995a, *JKAS*, 28, 169
 Lee, M. G. 1995b, *AJ*, 110, 1129
 Lee, M. G. et al. 1999, *AJ*, submitted
 Longmore, A. J., Hawarden, T. G., Webster, B. L., Goss, W. M., & Mebold, U. 1978, *MN*, 183, 97P
 Longmore, A. J., Hawarden, T. G., Goss, W. M., Mebold, U., & Webster, B. L. 1982, *MN*, 200, 325
 Lyo, A.-R., & Lee, M. G. 1997, *Jour. Korean Astro. Soc.*, 30, 27
 Mateo, M. 1998, *ARA&A*, 36, 435
 Miller, B. 1996, *AJ*, 112, 991
 Minniti, D., & Zijlstra, A. A. 1996, *ApJ*, 467, L13
 Puche, D., & Carignan, C. 1988, *AJ*, 95, 1025
 Salaris, M., & Cassisi, S. 1998, *MN*, 298, 166
 Sandage, A. 1971, in *Nuclei of Galaxies*, ed. D.J.K. O’Connell (Amsterdam, North-Holland), 601
 Schechter, P., Mateo, M., & Saha, A. 1993, *PASP*, 105, 1342
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 van den Bergh, S. 1994, *AJ*, 107, 1328




FIG. 1.— A grayscale map of the V-band CCD image of UKS 2323–326. North is at the top and east is to the left. The size of the field is $7'.5 \times 7'.5$. Regions labelled as C, I, and F represent, respectively, the central region ($r < 44''$), the intermediate region ($44'' < r < 77''$) and the control field region ($r < 77''$).




FIG. 2.— (a) $V-(B-V)$ diagram of the measured stars in the C-region plus I-region of UKS 2323-326. Filled circles and open circles represent the stars in the C-region and I-region, respectively. (b) $V-(B-V)$ diagram of the measured stars in the F-region.




FIG. 3.— (a) I –($V-I$) diagram of the measured stars in the C-region plus I-region of UKS 2323–326. Filled circles and open circles represent the stars in the C-region and I-region, respectively. (b) I –($V-I$) diagram of the measured stars in the F-region.




FIG. 4.— I -band luminosity function of the red giant branch stars in the C-region and I-region. The tip of the red giant branch is labelled as TRGB. The thick solid line, dotted line and dashed lines represent the luminosity functions of the I-region, C-region, and F-region, respectively. The contribution due to field stars was subtracted from the luminosity functions of the C-region and the I-region.




FIG. 5.— $I-(V-I)$ diagram of the measured stars in the I-region (open circles) and C-region (filled circles) of UKS 2323–326 in comparison with the red giant branches of Galactic globular clusters. The solid curved lines show, from left to right, the loci of the giant branches of M15, M2, and NGC 1851, the metallicities of which are $[Fe/H] = -2.17$, -1.58 and -1.29 dex, respectively. The mean errors for the magnitudes and colors are illustrated by the error bars at the right.




FIG. 6.— Surface photometry of UKS 2323-326. (a) Surface brightness profiles vs radius along the major axis. B , V , R and I magnitudes are represented by the open squares, filled circles, open triangles, and crosses. (b) Differential colors vs radius along the major axis.




FIG. 7.— Fits to the surface brightness profiles of UKS 2323–326. (a) King model fitting. Filled circles and open squares represent, respectively, V and B surface brightness profiles. (b) Exponential law fitting.




FIG. 8.— $V-(B-V)$ diagram of the measured stars in the C-region (filled circles) and I-region (open circles) of UKS 2323-326 in comparison with theoretical isochrones. The solid lines represent the Padova isochrones for the metallicity of $Z = 0.001$ ($[\text{Fe}/\text{H}] = -1.2$ dex) and ages of 10, 30, and 100 Myrs. The mean errors for the magnitudes and colors are illustrated by the error bars at the right.




FIG. 9.— V-band luminosity function of the bright main-sequence stars with $(B - V) < 0.4$ in UKS 2323–326. The dashed line represents a linear fit to the data for the range of $21.4 < V < 23$ mag.

TABLE 1
JOURNAL OF OBSERVATIONS FOR UKS 2323–326.

Filters	T_{exp}	Air mass	FWHM	U.T.(Start)
<i>B</i>	300 s	1.68	1.1''	1994 Oct 7 09:41
<i>V</i>	3×700 s	1.66	1.0''	1994 Oct 7 09:00
<i>R</i>	3×500 s	1.63	1.0''	1994 Oct 7 08:27
<i>I</i>	4×400 s	1.66	0.9''	1994 Oct 7 07:48

TABLE 2.
PHOTOMETRY OF THE BRIGHT STARS WITH $V < 22.5$ MAG IN UKS2323–326.

ID	X(px)	Y(px)	<i>V</i>	(<i>B</i> – <i>V</i>)	(<i>V</i> – <i>R</i>)	(<i>V</i> – <i>I</i>)	ID	X(px)	Y(px)	<i>V</i>	(<i>B</i> – <i>V</i>)	(<i>V</i> – <i>R</i>)	(<i>V</i> – <i>I</i>)
173	810.9	196.6	17.83	0.63	0.41	0.82	1013	532.0	626.7	21.43	0.12	–0.10	0.01
1659	177.2	805.2	18.01	0.56	0.36	0.69	1445	856.5	713.0	21.48	0.79	0.52	1.02
1858	268.6	921.5	18.09	0.65	0.48	0.89	185	805.8	203.7	21.60	1.17	–0.84	–0.90
1840	661.3	904.0	18.50	0.75	0.56	1.00	1251	445.4	664.5	21.79	–0.21	–0.10	0.12
1014	520.5	626.8	18.93	0.47	0.76	1.09	1153	924.3	646.8	21.86	1.30	–0.70	–0.01
132	212.2	163.2	19.03	1.03	0.75	1.40	612	121.4	526.7	21.88	1.39	0.91	1.93
1055	532.2	632.3	19.93	0.14	–0.03		997	452.5	624.0	21.88	0.05	0.25	0.37
332	217.6	329.1	20.00	0.87	0.62	1.12	999	510.9	624.3	21.89	–0.24	0.06	0.10
1899	205.9	959.4	20.35	1.13	0.83	1.51	1134	490.3	643.0	21.95	–0.05	–0.14	0.15
1614	270.6	775.3	20.38	0.94	0.72	1.28	225	179.7	234.9	22.01	0.50	0.27	0.51
47	583.7	58.8	20.42	0.76	0.41	0.83	1047	459.0	630.9	22.02	0.58	0.70	1.16
1002	522.6	624.7	20.42	0.23	0.04	0.27	980	461.8	621.8	22.04	–0.01	–0.15	0.17
956	508.3	619.5	20.49	1.37	0.81	1.38	885	462.8	606.8	22.04	0.32	0.34	0.71
1140	512.3	644.1	20.59	–0.49	–0.03	–0.06	930	750.6	615.1	22.11	0.72	–0.20	–0.22
1896	191.1	953.4	20.60	0.52	0.28	0.67	335	667.4	330.6	22.15	1.26	1.95	2.46
1116	508.8	640.8	20.63	0.05	–0.03	0.01	963	452.1	619.9	22.18	0.18	0.33	0.56
251	222.9	261.1	20.71	0.65	0.42	0.82	1483	473.6	725.3	22.19	0.02	–0.01	–0.07
1058	521.9	632.7	20.81	1.43	0.87		1863	555.2	928.1	22.20	0.02	0.25	0.39
821	489.3	593.8	20.92	1.24	1.00		968	506.1	620.1	22.20	0.00	–0.30	–0.24
96	813.8	127.9	21.03	0.34	0.18	0.56	900	512.0	609.4	22.24	–0.08	–0.09	–0.05
234	826.4	247.7	21.11	1.45	0.95	2.15	1691	359.6	818.7	22.25	0.52	0.20	0.84
174	740.0	198.1	21.18	0.47	0.35	0.67	1415	449.9	701.5	22.26	0.25	0.10	0.47
975	957.8	620.9	21.20	0.30	0.30	0.61	554	925.4	496.3	22.38	0.90	0.25	0.75
21	473.4	27.0	21.26	1.56	0.35	0.16	406	714.3	369.2	22.40	0.88	0.14	0.59
1280	166.7	669.7	21.30	0.23	0.17	0.35	388	820.7	359.4	22.47	0.03	0.41	0.38
404	932.9	368.8	21.34	0.61	0.29	0.70	1024	486.6	627.7	22.48	0.20	0.62	1.08
1490	477.1	726.3	21.38	0.51	0.75	1.07	1609	241.3	773.7	22.49	0.70	1.49	0.82

TABLE 3
BVRI SURFACE PHOTOMETRY OF UKS 2323–326.

$r_{\text{eff}} ["]$	μ_B	μ_V	μ_R	μ_I	$r_{\text{out}} ["]$	B	V	R	I
3.4	23.41	22.90	22.60	22.22	4.8	18.73	18.17	17.84	17.45
7.6	23.32	22.82	22.53	22.18	9.6	17.05	16.58	16.28	15.93
12.2	23.36	22.90	22.59	22.22	14.4	16.20	15.73	15.44	15.09
17.0	23.46	22.93	22.64	22.28	19.2	15.62	15.14	14.85	14.47
21.7	23.62	23.05	22.76	22.35	24.0	15.16	14.68	14.36	13.99
26.5	23.80	23.27	22.96	22.56	28.8	14.83	14.36	14.04	13.67
31.3	24.10	23.56	23.22	22.88	33.6	14.63	14.14	13.82	13.44
36.1	24.40	23.83	23.50	23.16	38.4	14.48	13.97	13.65	13.26
40.9	24.83	24.14	23.79	23.42	43.2	14.37	13.85	13.53	13.14
45.7	25.01	24.38	24.07	23.75	48.0	14.28	13.76	13.43	13.06
50.5	25.52	24.66	24.43	24.11	52.8	14.22	13.69	13.36	12.99
55.3	25.51	24.87	24.63	24.35	57.6	14.17	13.63	13.30	12.94
60.0	25.89	25.23	24.87	24.66	62.4	14.13	13.58	13.25	12.90
64.8	26.15	25.37	25.16	24.86	67.2	14.10	13.53	13.21	12.86
69.6	26.49	25.54	25.33	25.39	72.0	14.07	13.50	13.18	12.83
74.4	26.48	25.77	25.57	25.41	76.8	14.04	13.46	13.14	12.80
79.2	26.46	26.05	25.88	25.84	81.6	13.98	13.40	13.10	12.76

TABLE 4
BASIC INFORMATION OF UKS 2323–326.

Parameter	Information	Reference
$\alpha_{1950}, \delta_{1950}$	$23^h 23^m 47^s .6, -32^\circ 39' 57''$	1
l, b	11.86 deg, -70.86 deg	1
HI heliocentric radial velocity, $v_\odot$	62 ± 5 km s $^{-1}$	1
Foreground reddening, $E(B - V)$	0.014 mag	2
Distance	$(m - M)_0 = 26.59 \pm 0.12, d = 2.08 \pm 0.12$ Mpc	this study
Central surface brightness	$\mu_B(0) = 23.4, \mu_V(0) = 22.9$ mag arcsec $^{-2}$	this study
Core radius	32 arcsec = 320 pc	this study
Standard radius, r_{25}	46 arcsec = 460 pc	this study
Holmberg radius, r_H	70 arcsec = 700 pc	this study
Apparent total magnitude ($< r_H$)	$B = 14.07$ mag, $V = 13.50$ mag	this study
Absolute total magnitude	$M_B = -12.58$ mag, $M_V = -13.14$ mag	this study
RGB metallicity, [Fe/H]	-1.98 ± 0.17 dex	this study
HI Flux	15 ± 3 Jy km $^{-1}$	1
HI mass, M_{HI}	$1.5 \pm 0.3 \times 10^7 M_\odot$	this study

REFERENCES.— (1) Longmore et al. (1978); (2) Schlegel, Finkbeiner & Davis (1998).